

Patient-Specific Functional Scale

Source: Stratford P, Gill C, Westaway M, Binkley J. Assessing disability and change on individual patients: a report of a patient specific measure. *Physiotherapy Canada* 1995;47:258-63.

The Patient-Specific Functional Scale is designed to measure disability in people with an orthopaedic condition.

Scoring instructions

The patient is asked to identify activities which they find difficult to perform as a result of their condition. The clinician then shows the patient the scale, and asks the patient to give a score out of 10 for each activity.

Interpretation of scores

Total score = sum of the activity scores/number of activities

Minimum detectable change (90%CI) for average score = 2 points

Minimum detectable change (90%CI) for single activity score = 3 points



WholeHealth Networks, Inc. (WHN)

Patient-Specific Functional Scale

This questionnaire can be used to quantify activity limitation and measure functional outcome for patients with an orthopaedic condition.

Clinician to read and fill in. Complete at the end of the history and prior to physical examination.

Initial assessment													
with as a result of your								rtant activities that you are unable to do or are having difficulty problem. Today, are there any activities that you are unable t problem? (Clinician: show scale to patient					
	Issess	ed you	ı on (st	ate pr						ne that you ha			
0 0 1 Unable to perform activity	2 3 4 5 6 7					7	8	9	10 Able to perform activity at the same level as before injury or problem (Date and score)				
Activity				I	nitial								
1.													
2.													

Patient-specific activity scoring scheme (Point to one number):

3. 4. 5.

Additional Additional